

# THE DEMEYER-KANZAKI GALOIS EXTENSION AND ITS SKEW GROUP RING

GEORGE SZETO AND LIANYONG XUE

Bradley University, USA

ABSTRACT. Several characterizations are given for a ring  $B$  being a DeMeyer-Kanzaki Galois extension with Galois group  $G$  in terms of the skew group ring  $B * G$ . Consequently, the results of S. Ikehata on commutative Galois algebras are generalized.

## 1. INTRODUCTION

In [5], the class of commutative Galois algebras  $B$  with Galois group  $G$  was characterized in terms of the Azumaya skew group ring  $B * G$  over  $B^G$  and the  $H$ -separable skew group ring  $B * G$  of  $B$  respectively, where  $B^G = \{a \in B \mid g(a) = a \text{ for all } g \in G\}$ . In [3], a broader class of DeMeyer-Kanzaki Galois extensions  $B$  with Galois group  $G$  was investigated where  $B$  is called a DeMeyer-Kanzaki Galois extension with Galois group  $G$  if  $B$  is an Azumaya algebra over its center  $C$  and  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$ . Further generalizations to Azumaya Galois extensions and to Hopf Azumaya Galois extensions were also given (see [2], [7]). The purpose of the present paper is to generalize the characterizations of a commutative Galois algebra  $B$  in terms of the skew group ring  $B * G$  as given by S. Ikehata (see [5]). We shall show the following equivalent statements:

- (1)  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ .

---

*2000 Mathematics Subject Classification.* 16S35, 16W20

*Key Words and Phrases.* Galois extensions, DeMeyer-Kanzaki Galois extensions, commutative Galois algebras, Azumaya algebras,  $H$ -separable extensions, skew group rings.

This work was done under the support of a Caterpillar Fellowship at Bradley University. The authors would like to thank Caterpillar Inc. for the support.

- (2) The skew group ring  $B * G$  is an Azumaya  $C^G$ -algebra and  $C$  is a maximal commutative separable subalgebra of  $V_{B * G}(B^G)$ , the commutator subring of  $B^G$  in  $B * G$ , over  $C^G$ .
- (3) The skew group ring  $B * G$  is an  $H$ -separable extension of  $B$  (= the Harata separable),  $B$  is a separable algebra over  $C^G$ , and  $J_g = \{0\}$  for each  $g \neq 1$  in  $G$  where  $J_g = \{b \in B \mid bx = g(x)b \text{ for all } x \in B\}$  for each  $g \in G$ .
- (4)  $B$  is a separable  $C^G$ -algebra,  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and  $C \otimes_{C^G} (B * G) \cong M_n(B)$  where  $M_n(B)$  is the matrix ring of order  $n$  over  $B$  and  $n$  is the order of  $G$ .
- (5)  $B$  is a separable  $C^G$ -algebra,  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and  $C \otimes_{C^G} V_{B * G}(B^G) \cong M_n(C)$  where  $M_n(C)$  is the matrix ring of order  $n$  over  $C$  and  $n$  is the order of  $G$ .

## 2. BASIC DEFINITIONS AND NOTATIONS

Throughout,  $B$  will represent a ring with 1,  $C$  the center of  $B$ ,  $G$  a finite automorphism group of  $B$  of order  $n$  for some integer  $n$ ,  $B^G$  the set of elements fixed under each element in  $G$ , and  $J_g = \{b \in B \mid bx = g(x)b \text{ for all } x \in B\}$  for each  $g \in G$ . For a subring  $A$  of  $B$  with the same identity 1, we denote the commutator subring of  $A$  in  $B$  by  $V_B(A)$ . Following the definitions given in [10], we call  $B$  a separable extension of  $A$  if there exist  $\{a_i, b_i \text{ in } B, i = 1, 2, \dots, m \text{ for some integer } m\}$  such that  $\sum a_i b_i = 1$ , and  $\sum b a_i \otimes b_i = \sum a_i \otimes b_i b$  for all  $b$  in  $B$  where  $\otimes$  is over  $A$ . An Azumaya algebra is a separable extension of its center. A ring  $B$  is called an  $H$ -separable extension of  $A$  if  $B \otimes_A B$  is isomorphic to a direct summand of a finite direct sum of  $B$  as a  $B$ -bimodule.  $B$  is called a Galois extension of  $B^G$  with Galois group  $G$  if there exist elements  $\{a_i, b_i \text{ in } B, i = 1, 2, \dots, m \text{ for some integer } m\}$  such that  $\sum_{i=1}^m a_i g(b_i) = \delta_{1,g}$  for each  $g \in G$ . A Galois extension  $B$  with Galois group  $G$  is called an Azumaya Galois extension if  $B^G$  is an Azumaya algebra over  $C^G$  (see [2],[7]), and a DeMeyer-Kanzaki Galois extension if  $B$  is an Azumaya algebra over  $C$  which is a Galois algebra over  $C^G$  with Galois group induced by and isomorphic with  $G$  (see [3],[6]).

Let  $P$  be a finitely generated and projective module over a commutative ring  $R$ . Then for a prime ideal  $p$  of  $R$ ,  $P_p (= P \otimes_R R_p)$  is a free module over  $R_p (= \text{the local$

ring of  $R$  at  $p$ ), and the rank of  $P_p$  over  $R_p$  is the number of copies of  $R_p$  in  $P_p$ , that is,  $\text{rank}_{R_p}(P_p) = m$  for some integer  $m$ . It is known that the  $\text{rank}_R(P)$  is a continuous function ( $\text{rank}_R(P)(p) = m$ ) from  $\text{Spec}(R)$  to the set of nonnegative integers with the discrete topology (see [4, Corollary 4.11, page 31]). We shall use the  $\text{rank}_R(P)$ -function for a finitely generated and projective module  $P$  over a commutative ring  $R$ .

### 3. CHARACTERIZATIONS

In this section, keeping all notations as given in section 2, we shall generalize the characterizations of a commutative Galois algebra as given by S. Ikehata (see [5]) to a DeMeyer-Kanzaki Galois extension  $B$  with Galois group  $G$  in terms of the skew group ring  $B * G$ . We begin with an equivalent condition for a commutative Galois algebra  $C$  with Galois group  $G$ .

**THEOREM 3.1.** Let  $C$  be a commutative ring with a finite automorphism group  $G$ . Then,  $C$  is a commutative Galois algebra with Galois group  $G$  if and only if  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and  $C \otimes_{C^G} (C * G) \cong M_n(C)$ .

**PROOF.** ( $\implies$ ) By Corollary 1.3 on page 85 in [4],  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and that  $C \otimes_{C^G} (C * G) \cong M_n(C)$  is a consequence of Theorem 2 in [5].

( $\impliedby$ ) Since  $C \otimes_{C^G} (C * G) \cong M_n(C)$ ,  $C \otimes_{C^G} (C * G)$  is an Azumaya algebra over  $C$ . But  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule by hypothesis, so  $C * G$  is an Azumaya  $C^G$ -algebra (see [4, Corollary 1.10, page 45]). Hence  $C$  is a commutative Galois algebra with Galois group  $G$  (see [5, Theorem 2]).

Next we characterize a DeMeyer-Kanzaki Galois extension  $B$  in terms of the skew group ring  $B * G$ .

**THEOREM 3.2.** The following statements are equivalent:

- (1)  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ .
- (2) The skew group ring  $B * G$  is an Azumaya  $B^G$ -algebra and  $B$  is a maximal commutative separable subalgebra of  $V_{B * G}(B^G)$  over  $B^G$ .

- (3) The skew group ring  $B * G$  is an  $H$ -separable extension of  $B$ ,  $B$  is a separable algebra over  $C^G$ , and  $J_g = \{0\}$  for each  $g \neq 1$  in  $G$ .

PROOF. (1)  $\implies$  (2) Since  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ ,  $B \cong B^G \otimes_{C^G} C$  such that  $B^G$  is an Azumaya  $C^G$ -algebra (see [3, Lemma 2]). Hence  $B$  is an Azumaya Galois extension with Galois group  $G$ ; and so  $B * G$  is an Azumaya  $C^G$ -algebra (see [2, Theorem 1]). Moreover,  $C$  is a commutative Galois algebra with Galois group  $G$  by hypothesis, so  $C$  is a maximal commutative separable subalgebra of  $C * G$  over  $C^G$  (see [5, Theorem 2]). But  $V_{B * G}(B^G) = V_B(B^G) * G = C * G$ , so  $C$  is a maximal commutative separable subalgebra of  $V_{B * G}(B^G)$  over  $C^G$ .

(2)  $\implies$  (1) Since  $B * G$  is an Azumaya  $C^G$ -algebra,  $B$  is an Azumaya Galois extension with Galois group  $G$  (see [2, Theorem 1]). Hence  $V_B(B^G)$  is a Galois algebra over  $C^G$  with Galois group  $G$  (see [1, Theorem 2]). Thus  $V_B(B^G) * G \cong \text{Hom}_{C^G}(V_B(B^G), V_B(B^G))$ . But  $C$  is a maximal commutative separable subalgebra of  $V_B(B^G) * G (= V_{B * G}(B^G))$  over  $C^G$  by hypothesis, so by the proof of Theorem 5.5 on page 64 in [4],

$$C \otimes_{C^G} (V_B(B^G) * G) \cong \text{Hom}_C(V_B(B^G) * G, V_B(B^G) * G).$$

Then we have

$$\begin{aligned} \text{Hom}_C(V_B(B^G) * G, V_B(B^G) * G) &\cong C \otimes_{C^G} (V_B(B^G) * G) \\ &\cong C \otimes_{C^G} \text{Hom}_{C^G}(V_B(B^G), V_B(B^G)) \\ &\cong \text{Hom}_C(C \otimes_{C^G} V_B(B^G), C \otimes_{C^G} V_B(B^G)). \end{aligned}$$

Thus  $V_B(B^G) * G \cong (C \otimes_{C^G} V_B(B^G)) \otimes_C P$  as a  $C$ -module for some finitely generated and projective  $C$ -module  $P$  such that  $\text{rank}_C(P) = 1$ . Since the rank of a Galois algebra is the order of the Galois group, applying the rank function on both sides of the above isomorphism, we have that

$$\begin{aligned} \text{rank}_C(V_B(B^G)) \cdot n &= \text{rank}_C(V_B(B^G) * G) = \text{rank}_C(C \otimes_{C^G} V_B(B^G)) \\ &= \text{rank}_{C^G}(V_B(B^G)) = n. \end{aligned}$$

This implies that  $\text{rank}_C(V_B(B^G)) = 1$ . Noting that  $V_B(B^G)$  is an Azumaya  $C$ -algebra and a finitely generated projective  $C^G$ -module, we conclude that  $V_B(B^G) = C$ ; and so  $C$

is a Galois algebra over  $C^G$  with Galois group  $G$ . Consequently,  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$  because  $B$  is also an Azumaya  $C$ -algebra.

(1)  $\implies$  (3) Since  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ ,  $B \cong B^G \otimes_{C^G} C$  such that  $B^G$  is an Azumaya  $C^G$ -algebra and  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$  (see [3, Lemma 2]). Hence  $B * G$  is an  $H$ -separable extension of  $B$  (see [9, Lemma 3.1 and Theorem 3.2]) and  $B$  is a separable algebra over  $C^G$ . Noting that  $V_B(B^G) = C = J_1$  and that  $V_B(B^G) = \bigoplus \sum_{g \in G} J_g$  (see [6, Proposition 1]), we conclude that  $J_g = \{0\}$  for each  $g \neq 1$  in  $G$ .

(3)  $\implies$  (1) Since  $B$  is a separable algebra over  $C^G$ ,  $B$  is an Azumaya algebra over  $C$ . Next we claim that  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$ . In fact, since  $B * G$  is an  $H$ -separable extension of  $B$  by hypothesis and  $B$  is a direct summand of  $B * G$  as a left (or right)  $B$ -module,  $V_{B * G}(V_{B * G}(B)) = B$  (see [8, Proposition 1.2]). This implies that the center of  $B * G$  is  $C^G$ . Moreover,  $B$  is a separable algebra over  $C^G$ , so  $B * G$  is a separable algebra over  $C^G$  by the transitivity of separable extensions. Thus  $B * G$  is an Azumaya  $C^G$ -algebra; and so  $B$  is an Azumaya Galois extension with Galois group  $G$  (see [2, Theorem 1]). Therefore  $V_B(B^G)$  is a Galois algebra over  $C^G$  with Galois group induced by and isomorphic with  $G$  (see [1, Theorem 2]). But then, by Proposition 1 in [6],  $V_B(B^G) = \bigoplus \sum_{g \in G} J_g$ . Since  $J_g = \{0\}$  for each  $g \neq 1$  in  $G$  by hypothesis, so  $V_B(B^G) = J_1 = C$ . This proves that  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$ . Thus statement (1) holds.

By generalizing Theorem 3.1, we obtain another two characterizations of a DeMeyer-Kanzaki Galois extension.

**THEOREM 3.3.** The following statements are equivalent:

- (1)  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ .
- (2)  $B$  is a separable  $C^G$ -algebra,  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and  $C \otimes_{C^G} (B * G) \cong M_n(B)$ .
- (3)  $B$  is a separable  $C^G$ -algebra,  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule, and  $C \otimes_{C^G} V_{B * G}(B^G) \cong M_n(C)$ .

PROOF. (1)  $\implies$  (2) Since  $B$  is a DeMeyer-Kanzaki Galois extension of  $B^G$  with Galois group  $G$ ,  $B \cong B^G \otimes_{C^G} C$  where  $B^G$  is an Azumaya  $C^G$ -algebra and  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$  (see [3, Lemma 2]). Hence  $C^G$  is a direct summand of  $C$  as a  $C^G$ -submodule (see [4, Corollary 1.3, page 85]), and  $V_{B^G}(B^G) = C * G$  such that  $C \otimes_{C^G} (C * G) \cong M_n(C)$  (see [5, Theorem 2]); and so

$$\begin{aligned} C \otimes_{C^G} (B * G) &\cong C \otimes_{C^G} (B^G \otimes_{C^G} C * G) \cong C \otimes_{C^G} (C * G) \otimes_{C^G} B^G \\ &\cong M_n(C) \otimes_{C^G} B^G \cong M_n(B). \end{aligned}$$

(2)  $\implies$  (1) Since  $B$  is a separable  $C^G$ -algebra,  $B$  is an Azumaya algebra over  $C$ . Moreover,  $M_n(B) \cong B \otimes_C M_n(C)$ , so  $M_n(B)$  is an Azumaya  $C$ -algebra. By hypothesis,  $C \otimes_{C^G} (B * G) \cong M_n(B)$ , so  $C \otimes_{C^G} (B * G)$  is an Azumaya algebra over  $C$ . But  $C$  contains  $C^G$  as a direct summand as a  $C^G$ -submodule by hypothesis, so  $B * G$  is an Azumaya  $C^G$ -algebra (see [4, Corollary 1.10, page 45]). Hence  $B$  is an Azumaya Galois extension with Galois group  $G$  (see [2, Theorem 1]). Thus  $V_B(B^G)$  is a Galois algebra over  $C^G$  with Galois group  $G$  (see [1, Theorem 2]). Therefore both  $B$  and  $B^G \cdot V_B(B^G)$  are Galois extensions of  $B^G$  with Galois group  $G$  such that  $B^G \cdot V_B(B^G) \subset B$ . This implies that  $B = B^G \cdot V_B(B^G)$  such that  $V_B(B^G)$  is a Galois algebra over  $C^G$  with Galois group  $G$ ; and so  $V_B(B^G)$  an Azumaya  $C$ -algebra and both  $V_B(B^G)$  and  $C$  are finitely generated projective modules over  $C^G$ .

Next we claim that  $V_B(B^G) = C$ . In fact, since  $C \otimes_{C^G} (B * G) \cong M_n(B)$ ,  $\text{rank}_{C^G}(B * G) = \text{rank}_C(M_n(B))$ . This implies that  $\text{rank}_{C^G}(C) \cdot \text{rank}_C(B) \cdot n = \text{rank}_C(B) \cdot n^2$ . Thus  $\text{rank}_{C^G}(C) = n$ . But  $V_B(B^G)$  is a Galois algebra over  $C^G$  with Galois group  $G$ , so  $\text{rank}_{C^G}(V_B(B^G)) = n$ . Therefore  $\text{rank}_{C^G}(V_B(B^G)) = n = \text{rank}_{C^G}(C)$ . Noting that  $V_B(B^G)$  is an Azumaya  $C$ -algebra and a finitely generated projective  $C^G$ -module, we conclude that  $V_B(B^G) = C$ ; and so  $C$  is a Galois algebra with Galois group induced by and isomorphic with  $G$ . Consequently,  $B$  is a DeMeyer-Kanzaki Galois extension with Galois group  $G$ .

(1)  $\iff$  (3) The proof is similar to (1)  $\iff$  (2).

## REFERENCES

- [1] R. Alfaro and G. Szeto, Skew Group Rings which are Azumaya, *Comm. in Algebra*, 23(6)(1995), 2255-2261.
- [2] R. Alfaro and G. Szeto, On Galois Extensions of an Azumaya Algebra, *Comm. in Algebra*, 25(6)(1997), 1873-1882.
- [3] F.R. DeMeyer, Some Notes on the General Galois Theory of Rings, *Osaka J. Math.*, 2(1965), 117-127.
- [4] F. R. DeMeyer and E. Ingraham, *Separable algebras over commutative rings*, Volume 181. Springer Verlag, Berlin, Heidelberg, New York, 1971.
- [5] S. Ikehata, Note on Azumaya Algebras and  $H$ -Separable Extensions, *Math. J. Okayama Univ.*, 23(1981), 17-18.
- [6] T. Kanzaki, On Galois algebra over a commutative ring, *Osaka J. Math.*, 2(1965), 309-317.
- [7] M. Ouyang, Azumaya Extensions and Galois Correspondence, *Algebra Colloquium*, 7(1) (2000), 43-57.
- [8] K. Sugano, Note on Semisimple Extensions and Separable Extensions, *Osaka J. Math.*, 4(1967) 265-270.
- [9] G. Szeto and L. Xue, On Characterizations of a Center Galois Extension, *International Journal of Mathematics and Mathematical Sciences*, Vol. 23, No. 11(2000), 753-758.
- [10] G. Szeto and L. Xue, The structure of Galois algebras, *Journal of Algebra*, 237(1)(2001), 238-246.

Department of Mathematics  
Bradley University  
Peoria, Illinois 61625, USA

E-mail address: szeto@hilltop.bradley.edu

E-mail address: lxue@hilltop.bradley.edu